

ONGLET 3 :

DAVIDOU Ludivine, Jean-Christophe CRACHEREAU - Mise en évidence de l'impact qualitatif de la présence de mildiou (*Plasmopara viticola*) en faciès Rot Brun sur les vins rouges de Bordeaux et détermination de seuils de tolérance revue agriculture et territoires chambre d'agriculture de la gironde, 2012

http://www.vinopole.com/uploads/tx_vinoexperimentation/Gout_Mildiou.pdf

Marchand S., Almy J., de Revel G.

The cysteine reaction with diacetyl under wine- like conditions: Proposed mechanisms for mixed

origins of 2-methylthiazole, 2-methyl-3- thiazoline, 2-methylthiazolidine, and 2,4,5-trimethyloxazole,

Journal of Food Science 76 (6) : C861-C86, 2011.

Nikolantonaki M., Darriet P.

Identification of ethyl 2-sulfanylacetate as an important off-odor compound in white wines,

Journal of Agricultural and Food Chemistry 59 (18) : 10191-10199, 2011.

Pons M., Dauphin B., La Guerche S., Pons A., Lavigne-Cruege V., Shinkaruk S., Bunner, D., Darriet P. **Identification of impact odorants contributing to fresh mushroom off-flavor in wines: Incidence of their reactivity with nitrogen compounds on the decrease of the olfactory defect,** *Journal of Agricultural and Food Chemistry* 59 (7) : 3264-3272

Tempere S., Cuzange E., Malak J., Bougeant J.C., De Revel G., Sicard G.

The training level of experts influences their detection thresholds for key wine compounds,

Chemosensory Perception 4 (3) : 99-115, 2011

Tempere S., Cuzange E., Malak J., Bougeant J.C., De Revel G., Sicard G.

Cartographie des capacités olfactives des professionnels du vin

Rev. Des Œnologues, janvier 2012, n°142